



Pr **Tegsedi**[®]
(inotersen) *injection*
284 mg/1.5 mL

 sobi

 AKCEA CONNECT
SUPPORT PROGRAM

TEGSEDI[®] and AKCEA[®] CONNECT are registered trademarks of Akcea Therapeutics Canada, Inc. Discovered and developed by Ionis Pharmaceuticals. ©2021 Akcea Therapeutics Canada, Inc. All rights reserved. PP-11844_07/21

 MEMBER OF
INNOVATIVE MEDICINES CANADA

TRAVEL CHECKLIST



AKCEA CONNECT
SUPPORT PROGRAM

AKCEA[®] CONNECT

SUPPORT PROGRAM

Akcea[®] Connect is here to help connect you to resources and provide additional support for your P^rTEGSEDI[®] treatment. This includes helping to coordinate services through specialty clinic networks while you may be traveling.

TEGSEDI is used for the treatment of stage 1 or stage 2 polyneuropathy in adult patients with hereditary transthyretin amyloidosis (hATTR).

Travel checklist

Here are a few things to consider before you travel:

- Contact your Nurse Case Manager to see how Akcea Connect can support you during this period. Keep your Nurse Case Manager's phone number with you while you travel:



1-833-327-0723

- Coordinate with your healthcare provider before departure and inform them when you will be traveling:



From DD MM YYYY to DD MM YYYY

- Ask your healthcare provider to outline any necessary administration support and laboratory monitoring and record it below:



Note: _____

- Verify that you will be able to store TEGSEDI in the original container at the required temperature (2 °C to 8 °C) away from light. TEGSEDI may be stored unrefrigerated (2 °C to 30 °C) in the original container for up to 6 weeks; if not used within the 6 weeks, throw TEGSEDI away.



- Storage requirements

- Confirm that you will have enough of the additional necessary supplies for injection, including:



- Alcohol wipe
- Gauze pad or cotton ball
- Sharps container

For more information on the Akcea Connect Patient Support Program, please contact us by:



1-833-327-0723



support@akceaconnect.ca

Or visit our website at **akceaconnect.ca**


Pr **Tegsedi**[®]
(inotersen) injection
284 mg/15 mL